

*Suppl I,  
JP XVII  
(2017)*

*Suppl I,  
JP XVII  
(2017)*

*Suppl I,  
JP XVII  
(2017)*

### Hangeshashinto Extract

半夏瀉心湯エキス

**Change the Origin/limits of content, Identification (2), (3), (5) and Assay (2) as follows:**

Hangeshashinto Extract contains not less than 70 mg and not more than 210 mg (for preparation prescribed 2.5 g of Scutellaria Root) or not less than 80 mg and not more than 240 mg (for preparation prescribed 3 g of Scutellaria Root) of baicalin ( $C_{21}H_{18}O_{11}$ : 446.36), not less than 18 mg and not more than 54 mg (for preparation prescribed 2.5 g of Glycyrrhiza) or not less than 20 mg and not more than 60 mg (for preparation prescribed 3 g of Glycyrrhiza) of glycyrrhizic acid ( $C_{42}H_{62}O_{16}$ : 822.93), and not less than 7 mg and not more than 21 mg of berberine [as berberine chloride ( $C_{20}H_{18}ClNO_4$ : 371.81)], per extract prepared with the amount specified in the Method of preparation.

#### Identification

(2) For preparation prescribed Processed Ginger— Shake 1.0 g of the dry extract (or 3.0 g of the viscous extract) with 10 mL of water, add 25 mL of diethyl ether, and shake. Separate the diethyl ether layer, evaporate the layer under reduced pressure, add 2 mL of diethyl ether to the residue, and use this solution as the sample solution. Separately, dissolve 1 mg of [6]-shogaol for thin-layer chroma-

*Suppl I,  
JP XVII  
(2017)*

tography in 1 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 20  $\mu\text{L}$  of the sample solution and 1  $\mu\text{L}$  of the standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of cyclohexane and ethyl acetate (2:1) to a distance of about 7 cm, and air-dry the plate. Spray evenly 4-dimethylaminobenzaldehyde TS for spraying on the plate, heat the plate at 105°C for 5 minutes, allow to cool, and spray water: one of the several spots obtained from the sample solution has the same color tone and *R<sub>f</sub>* value with the blue-green to grayish green spot from the standard solution (Processed Ginger).

(3) For preparation prescribed Ginger—Shake 1.0 g of the dry extract (or 3.0 g of the viscous extract) with 10 mL of water, add 25 mL of diethyl ether, and shake. Separate the diethyl ether layer, evaporate the layer under reduced pressure, add 2 mL of diethyl ether to the residue, and use this solution as the sample solution. Separately, dissolve 1 mg of [6]-gingerol for thin-layer chromatography in 1 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 10  $\mu\text{L}$  of the sample solution and 5  $\mu\text{L}$  of the standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate and hexane (1:1) to a distance of about 7 cm, and air-dry the plate. Spray evenly 4-dimethylaminobenzaldehyde TS for spraying on the plate, heat the plate at 105°C for 5 minutes, allow to cool, and spray water: one of the several spots obtained from the sample solution has the same color tone and *R<sub>f</sub>* value with the blue-green to grayish green spot from the standard solution (Ginger).

(5) Shake 1.0 g of the dry extract (or 3.0 g of the viscous extract) with 10 mL of water, add 5 mL of 1-butanol, shake, centrifuge, and use the supernatant liquid as the sample solution. Separately, dissolve 1 mg of liquiritin for thin-layer chromatography in 1 mL of methanol, and use this solution as the standard solution. Perform the test with these solutions as directed under Thin-layer Chromatography <2.03>. Spot 1  $\mu\text{L}$  each of the sample solution and standard solution on a plate of silica gel for thin-layer chromatography. Develop the plate with a mixture of ethyl acetate, methanol and water (20:3:2) to a distance of about 7 cm, and air-dry the plate. Spray evenly dilute sulfuric acid on the plate, heat the plate at 105°C for 5 minutes, and examine under ultraviolet light (main wavelength: 365 nm): one of the several spots obtained from the sample solution has the same color tone and *R<sub>f</sub>* value with the yellow-green fluorescent spot from the standard solution (Glycyrrhiza).

#### Assay

(2) Glycyrrhizic acid—Perform the test according to the following i) or ii).

i) Weigh accurately about 0.5 g of the dry extract (or an amount of the viscous extract, equivalent to about 0.5 g of the dried substance), add exactly 50 mL of diluted methanol (1 in 2), shake for 15 minutes, filter, and use the filtrate as

the sample solution. Separately, weigh accurately about 10 mg of Glycyrrhizic Acid RS (separately determine the water <2.48> by coulometric titration, using 10 mg), dissolve in diluted methanol (1 in 2) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu\text{L}$  each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas, *A<sub>T</sub>* and *A<sub>S</sub>*, of glycyrrhizic acid in each solution.

$$\begin{aligned} &\text{Amount (mg) of glycyrrhizic acid (C}_{42}\text{H}_{62}\text{O}_{16}) \\ &= M_S \times A_T / A_S \times 1/2 \end{aligned}$$

*M<sub>S</sub>*: Amount (mg) of Glycyrrhizic Acid RS taken, calculated on the anhydrous basis

#### Operating conditions—

Detector: An ultraviolet absorption photometer (wavelength: 254 nm).

Column: A stainless steel column 4.6 mm in inside diameter and 15 cm in length, packed with octadecylsilanized silica gel for liquid chromatography (5  $\mu\text{m}$  in particle diameter).

Column temperature: A constant temperature of about 40°C.

Mobile phase: Dissolve 3.85 g of ammonium acetate in 720 mL of water, and add 5 mL of acetic acid (100) and 280 mL of acetonitrile.

Flow rate: 1.0 mL per minute (the retention time of glycyrrhizic acid is about 15 minutes).

#### System suitability—

System performance: Dissolve 5 mg of monoammonium glycyrrhizinate for resolution check in 20 mL of dilute ethanol. When the procedure is run with 10  $\mu\text{L}$  of this solution under the above operating conditions, the resolution between the peak having the relative retention time of about 0.9 to glycyrrhizic acid and the peak of glycyrrhizic acid is not less than 1.5. Dissolve 1 mg of baicalein for resolution check in 50 mL of methanol. To 2 mL of this solution add 2 mL of the standard solution. When the procedure is run with 10  $\mu\text{L}$  of this solution under the above operating conditions, the resolution between the peaks of glycyrrhizic acid and baicalein is not less than 1.5.

System repeatability: When the test is repeated 6 times with 10  $\mu\text{L}$  of the standard solution under the above operating conditions, the relative standard deviation of the peak area of glycyrrhizic acid is not more than 1.5%.

ii) Weigh accurately about 0.5 g of the dry extract (or an amount of the viscous extract, equivalent to about 0.5 g of the dried substance), add 20 mL of ethyl acetate and 10 mL of water, and shake for 10 minutes. After centrifugation, remove the upper layer, add 20 mL of ethyl acetate, proceed in the same manner as described above, and remove the upper layer. To the resultant aqueous layer add 10 mL of methanol, shake for 30 minutes, centrifuge, and take the supernatant liquid. To the residue add 20 mL of diluted methanol (1 in 2), shake for 5 minutes, centrifuge, and take the supernatant liquid. Combine these supernatant liquids, add diluted methanol (1 in 2) to make exactly 50 mL, and use

this solution as the sample solution. Separately, weigh accurately about 10 mg of Glycyrrhizic Acid RS (separately determine the water <2.48> by coulometric titration, using 10 mg), dissolve in diluted methanol (1 in 2) to make exactly 100 mL, and use this solution as the standard solution. Perform the test with exactly 10  $\mu$ L each of the sample solution and standard solution as directed under Liquid Chromatography <2.01> according to the following conditions, and determine the peak areas,  $A_T$  and  $A_S$ , of glycyrrhizic acid in each solution.

$$\begin{aligned} & \text{Amount (mg) of glycyrrhizic acid (C}_{42}\text{H}_{62}\text{O}_{16}\text{)} \\ & = M_S \times A_T / A_S \times 1/2 \end{aligned}$$

$M_S$ : Amount (mg) of Glycyrrhizic Acid RS taken, calculated on the anhydrous basis

*Operating conditions—*

Proceed as directed in the operating conditions in i).

*System suitability—*

System repeatability: Proceed as directed in the system suitability in i).

System performance: Dissolve 5 mg of monoammonium glycyrrhizinate for resolution check in 20 mL of dilute ethanol. When the procedure is run with 10  $\mu$ L of this solution under the above operating conditions, the resolution between the peak having the relative retention time of about 0.9 to glycyrrhizic acid and the peak of glycyrrhizic acid is not less than 1.5.

*Suppl I,  
JP XVII  
(2017)*

*Suppl I,  
JP XVII  
(2017)*

*Suppl I,  
JP XVII  
(2017)*

*Suppl I,  
JP XVII  
(2017)*